

The following instructions should be followed closely except when overruled by specific procedural instructions.

Two Weeks Prior to Surgery

- NO ASPIRIN (*or medicines that contain aspirin*) since it interferes with normal blood clotting.
- NO IBUPROFEN (*or medicines contain ibuprofen*) as it interferes with blood clotting.
- Please DISCONTINUE ALL HERBAL VITAMINS OR MEDICATIONS as many have side effects that could complicate a surgical procedure by inhibiting blood clotting, affecting blood pressure, or interfering with anesthetics.
- Please DISCONTINUE ALL DIET PILLS whether prescription, over-the-counter or herbal as many will interfere with anesthesia and can cause cardiovascular concerns.
- NO SMOKING OR SECOND HAND SMOKE EXPOSURE because nicotine reduces blood flow to the skin and can cause significant complications during healing for two weeks prior to surgery and for four weeks following procedure.
- You may take Tylenol or generic forms of this drug. These do **not** interfere with blood clotting or healing.
- Contact our office should your medical history have changed since your consultation (i.e. any new medications, medical concerns, other medical investigations being done by any other Physicians in your care).

• One Week Prior to Surgery

- DO NOT take or drink any alcohol or drugs for one week prior to surgery and one week after surgery as these can interfere with anesthesia and affect blood clotting.
- If your skin tolerates, use a germ-inhibiting soap for bathing, such as Dial, Safeguard, or Lever 2000 for at least the week before surgery.
- DO report any signs of cold, infection, boils, or pustules (or new medical concerns) appearing before surgery.
- DO NOT take any cough or cold medications without permission.
- DO arrange for a responsible adult to drive you to and from the facility on the day of surgery, since you will not be allowed to leave on your own.
- DO arrange for a responsible individual to spend the first 24 hours after surgery with you, since you CANNOT be left alone.

• Night Before Surgery & Morning of Surgery

- DO NOT eat or drink anything (not even water) after midnight the night before your surgery. Also, no gum, candy, mints or coffee the morning of surgery. Do not sneak anything as this may endanger you.
- If you are on regular medications, please clear these with us.
- DO take a thorough shower with your germ-inhibiting soap the night before and the morning of surgery. Shampoo your hair the morning of surgery. This is to decrease the bacteria on the skin and thereby decrease the risk of infection.
- DO NOT apply any of the following to your skin, hair or face the morning of surgery: makeup, creams, lotions, hair gels, sprays, perfumes, powder, or deodorant. Using any of these products will add bacteria to the skin and increase the risk of infection.

- You may brush your teeth the morning of surgery but do not drink anything.
- DO NOT wear contacts to surgery. If you do wear glasses, bring your eyeglass case.
- DO wear comfortable, loose-fitting clothes that do not have to be put on over your head. The best thing to wear home is a button-up top and pull on pants. You will want easy-to-slip-on flat shoes.
- DO NOT bring any valuables or wear any jewelry (no rings, earrings, chains, toe rings, other metal piercings or watches). We will need to tape wedding rings if worn.
- You must have an adult drive for you – to and from surgery. Please note that a cab or bus driver will not be allowed to take you home after surgery. On arrival, be sure we know your driver's name, phone numbers, and how we will be able to reach them.
- If you are not recovering at home, it is very important that we have the number where you will be after surgery.

The following instructions should be followed closely except when overruled by specific procedural instructions. You must follow the instructions as indicated for your specific surgery. Notify Dr. Barr of any unusual changes in your condition and feel free to call the office with any questions.

- You **MUST HAVE AN ADULT DRIVE YOU** home from the facility. You will **not** be allowed to drive yourself or use public transportation.
- After surgery you **MUST HAVE A RESPONSIBLE ADULT STAY WITH YOU** a minimum of 24 hours. You **CANNOT** be left alone. The 24 hours begin when you are discharged from the office or hospital. Have everything ready at home **PRIOR** to surgery. Make arrangements for someone to stay with you. Let the person or persons know you cannot be left alone. This is important because of the danger of falling and you may lose the concept of time for the day and overmedicate yourself.
- The effects of anaesthesia can persist for 24 hours. You must exercise extreme caution before engaging in any activity that could be harmful to yourself or others.
- **DRINK** fluids to help rid the body of the drugs used in surgery. If you have straws in the house you will tend to drink more fluids the first few days after surgery.
- Diet may be as tolerated. Eating foods that are bland and soft for the first day or so – foods like after you have had the flu – may be best tolerated. You must eat more than crackers and juice, otherwise you will continue to feel weak and will not heal as well. **REMEMBER** to take the medications with a little something to eat or you will get sick to your stomach.
- Please avoid the use of alcoholic beverages for the first 24 hours (it dilates blood vessels and can cause unwanted bleeding) and as long as pain medications are being used (dangerous combination).
- Take only medications that have been prescribed by Dr. Barr for your postoperative care and take them according to the instruction on the bottle. Your pain medication may make you feel “spacey”; therefore, have someone else give you your medications according to the proper time intervals.
- If you experience any generalized itching, rash, wheezing or tightness in the throat, stop taking all medications and call the office immediately, as this may be a sign of a drug allergy.
- You can expect moderate discomfort, which should be helped by the pain medications. The greatest discomfort is usually during the first 24 hours. Thereafter, you will find that you require less pain medication.
- Call 523-4900 extension “0” (*toll-free 1-877-523-4900*) if you have: **SEVERE PAIN** not responding to pain medication; Swelling that is greater on one side than the other; incisions that are **RED OR FEVERISH**; a **FEVER**; or if any other questions or problems arise.
- Keep any **DRESSINGS ON, CLEAN AND DRY**. Do not remove them until instructed to do so. There may be some bloody drainage on the dressings. If you have excessive bleeding or the bandages are too tight, call the office immediately.
- After surgery it is important to have a bowel movement within a day or two. If you do not, you may take over-the-counter laxatives to encourage your bowels to move.
- You may note that when you stand up very quickly after lying down, that you feel slightly lightheaded. If this is a problem for you, sit with your feet off the edge of the bed for a few moments prior to standing up. You may also take a multi-vitamin with Iron once or twice daily for the next six to eight weeks.

POST-OPERATIVE INSTRUCTIONS - GENERAL

- Minimal activity for the first 48 hours. No house cleaning, furniture rearranging, etc. Relax, be pampered, and let your body heal. The less energy you use on doing things, the more energy your body can focus on healing.
- Limit "light" activities (under 20 pounds) i.e. lifting, pulling or pushing for **at least 10 days**. Return to heavier duties should be cleared by your surgeon (normally you will require 4 to 6 weeks to return to heavy duties).
- Position after surgery is different with different types of surgery. If your surgery is from the waist up we ask that the head of the bed be elevated 45 degrees. This requires a pillow under the small of your back, two pillows under your shoulders and head, and if you have a pillow under each elbow you will relax and stay in position.
- You are requested to remain within a reasonable traveling distance of the office for approximately ten days after surgery.
- Once cleared to shower you may do so every day. Please do not use the bathtub for 2 weeks.
- **NO SMOKING** for the first 4 weeks after specified surgeries (*this is a MUST for the following surgeries: Abdominoplasty, Rhytidectomy, BrowLift, Mastopexy, ThighLift, Brachioplasty, and Otoplasty*). Any cheating will delay healing.
- You may drive once you are off the pain pills, and when you experience no pain with this activity (as you need to be able to react quickly).
- All surgeries involve some scarring, which can take up to a year to fade. No matter how small they may be, we still want them to heal as well as they are able. Exposing red scars to the sun can cause permanent discoloration. A good sunscreen (SPF 15 or higher) can help and will protect the surrounding tissues that might not feel a sunburn developing while the nerves are healing. Sunlight can even reach scars under a swimsuit, so take adequate precautions.
- **DO NOT** use a hot tub for 4 weeks.
- **AVOID** sports or strenuous activities for 4 to 6 weeks or until our office gives you clearance during your post-operative visits. This is to avoid any unnecessary complications (bleeding, bruising, swelling).
- You may return to work when you feel able and are cleared to do so by our office.
- Feel free to call upon us at anytime. We want you to be as comfortable as possible during your healing period!